

# Exercise Guidelines for Cancer Survivors: Consensus Statement from International Multidisciplinary Roundtable

KRISTIN L. CAMPBELL<sup>1</sup>, KERRI M. WINTERS-STONE<sup>2</sup>, JOACHIM WISKEMANN<sup>3</sup>, ANNE M. MAY<sup>4</sup>, ANNA L. SCHWARTZ<sup>5</sup>, KERRY S. COURNEYA<sup>6</sup>, DAVID S. ZUCKER<sup>7</sup>, CHARLES E. MATTHEWS<sup>8</sup>, JENNIFER A. LIGIBEL<sup>9</sup>, LYNN H. GERBER<sup>10,11</sup>, G. STEPHEN MORRIS<sup>12</sup>, ALPA V. PATEL<sup>13</sup>, TRISHA F. HUE<sup>14</sup>, FRANK M. PERNA<sup>15</sup>, and KATHRYN H. SCHMITZ<sup>16</sup>

<sup>1</sup>Department of Physical Therapy, Faculty of Medicine, University of British Columbia, Vancouver, CANADA; <sup>2</sup>School of Nursing and Knight Cancer Institute, Oregon Health Sciences University, Portland, OR; <sup>3</sup>Division of Medical Oncology, National Center for Tumor Diseases (NCT) and Heidelberg University Clinic, Heidelberg, GERMANY; <sup>4</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, THE NETHERLANDS; <sup>5</sup>School of Nursing, Northern Arizona University, Flagstaff, AZ; <sup>6</sup>Faculty of Kinesiology, Sport, and Recreation, University of Alberta, Edmonton, CANADA; <sup>7</sup>Cancer Rehabilitation Medicine Services, Swedish Cancer Institute, Swedish Health Services, Seattle, WA; <sup>8</sup>Metabolic Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD; <sup>9</sup>Havard Medical School, Boston, MA; <sup>10</sup>Department of Medicine, Inova Fairfax Medical Campus, Falls Church, VA; <sup>11</sup>Center for the Study of Chronic Illness and Disability, George Mason University, Fairfax, VA; <sup>12</sup>Physical Therapy, Wingate University, Wingate, NC; <sup>13</sup>Epidemiology Research, American Cancer Society, Atlanta, GA; <sup>14</sup>Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, CA; <sup>15</sup>Division of Cancer Control and Population Sciences, Behavioral Research Program, Health Behaviors Research Branch, National Cancer Institute, Rockville, MD; and <sup>16</sup>Public Health Science, Penn State Cancer Institute, Penn State College of Medicine, Hershey, PA

## ABSTRACT

CAMPBELL, K. L., K. M. WINTERS-STONE, J. WISKEMANN, A. M. MAY, A. L. SCHWARTZ, K. S. COURNEYA, D. S. ZUCKER, C. E. MATTHEWS, J. A. LIGIBEL, L. H. GERBER, G. S. MORRIS, A. V. PATEL, T. F. HUE, F. M. PERNA, and K. H. SCHMITZ. Exercise Guidelines for Cancer Survivors: Consensus Statement from International Multidisciplinary Roundtable. *Med. Sci. Sports Exerc.*, Vol. 51, No. 11, pp. 2375–2390, 2019. **Purpose:** The number of cancer survivors worldwide is growing, with over 15.5 million cancer survivors in the United States alone—a figure expected to double in the coming decades. Cancer survivors face unique health challenges as a result of their cancer diagnosis and the impact of treatments on their physical and mental well-being. For example, cancer survivors often experience declines in physical functioning and quality of life while facing an increased risk of cancer recurrence and all-cause mortality compared with persons without cancer. The 2010 American College of Sports Medicine Roundtable was among the first reports to conclude that cancer survivors could safely engage in enough exercise training to improve physical fitness and restore physical functioning, enhance quality of life, and mitigate cancer-related fatigue. **Methods:** A second Roundtable was convened in 2018 to advance exercise recommendations beyond public health guidelines and toward prescriptive programs specific to cancer type, treatments, and/or outcomes. **Results:** Overall findings retained the conclusions that exercise training and testing were generally safe for cancer survivors and that every survivor should “avoid inactivity.” Enough evidence was available to conclude that specific doses of aerobic, combined aerobic plus resistance training, and/or resistance training could improve common cancer-related health outcomes, including anxiety, depressive symptoms, fatigue, physical

Address for correspondence: Kristin L. Campbell, B.Sc. P.T., M.Sc., Ph.D., F.A.C.S.M., Department of Physical Therapy, University of British Columbia, 212, 2177 Wesbrook Mall, Vancouver, British Columbia, Canada V6T 1Z3; E-mail: Kristin.campbell@ubc.ca.

K.L.C. and K.W.-S. are the co-first authors.

Submitted for publication June 2019.

Accepted for publication June 2019.

0195-9131/19/5111-2375/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2019 by the American College of Sports Medicine

DOI: 10.1249/MSS.0000000000002116

functioning, and health-related quality of life. Implications for other outcomes, such as peripheral neuropathy and cognitive functioning, remain uncertain. **Conclusions:** The proposed recommendations should serve as a guide for the fitness and health care professional working with cancer survivors. More research is needed to fill remaining gaps in knowledge to better serve cancer survivors, as well as fitness and health care professionals, to improve clinical practice. **Key Words:** CANCER SURVIVORSHIP, EXERCISE PRESCRIPTION, SAFETY

In the last decade, the United States has seen a 27% decline in cancer deaths due to early detection and improved treatments for cancer. In turn, the number of cancer survivors is growing, with over 15.5 million cancer survivors in the United States—a figure that is expected to double by 2040 (1). Improved prognosis has created a growing need to address the unique health issues facing cancer survivors that result from the disease, its treatment, and related comorbid conditions. For example, the symptom of fatigue can persist in 25% of cancer survivors many years after their treatment has ended and contributes to difficulty returning to work, independent living, and poor quality of life (2). Furthermore, risk of developing heart disease may be elevated by some cancer treatments, and cardiovascular mortality is emerging as a major competing cause of death in cancer survivors along with cancer recurrence (3,4). Cancer is also a disease strongly linked with aging, and almost half of survivors are older than 70 yr (5). The adverse synergistic effects of age, cancer treatment, and related sequelae increase the total burden of cancer. Historically, clinicians advised cancer patients to rest and to avoid physical activity, but early exercise research in the 1990s and 2000s challenged this advice.

In 2010, the American College of Sports Medicine convened a Roundtable meeting composed of a team of clinical and research experts in the field of cancer and exercise to develop the first set of exercise guidelines for cancer survivors (6). Drawing on studies mainly in breast and prostate cancer survivors, the key findings from this review were that exercise training was generally safe and well tolerated during and after cancer treatment and could elicit improvements in some health outcomes. There was sufficient evidence to conclude that exercise could improve physical fitness, physical functioning,

quality of life, and cancer-related fatigue. However, given the limited number of rigorously designed studies at that time, there was insufficient evidence to inform specific exercise prescriptions for any of these outcomes nor by cancer site or treatment type. Thus, the 2010 Roundtable recommendations largely followed the 2008 Physical Activity Guidelines for adults with chronic conditions to aim for at least 150 min·wk<sup>-1</sup> of aerobic activity, two or more days a week of resistance training, and daily stretching of major muscle groups when possible, with specific exercise testing and program modifications based on health status and cancer and treatment-related side effects (7). At a minimum, survivors were urged to “avoid inactivity” and be as physically active as possible (6).

The 2010 American College of Sports Medicine (ACSM) recommendations came with the expectation that they would be updated as the evidence grew and indeed, since that landmark publication, the number of randomized controlled exercise trials in cancer survivors has increased by 281% (PubMed Search completed March 2018) to over 2500 published randomized controlled trials. Moreover, an increasing number of calls for the integration of exercise into clinical cancer care have since been issued (8–11). Thus, in 2018, the ACSM International Multidisciplinary Roundtable on Physical Activity and Cancer Prevention and Control was convened to bring together an international group of exercise and rehabilitation professionals and organizations with the goal to update recommendations based on current evidence. The Roundtable meeting took place on March 12 to 13, 2018, in San Francisco, CA, with 40 representatives from 20 organizations across the world who came together to sponsor and attend this meeting (Table 1). Roundtable members were invited to participate based on their clinical and scientific

TABLE 1. Professional and stakeholder organizations involved in ACSM International Multidisciplinary Roundtable on Exercise and Cancer

Participating Organizations	Partner Organization	Official Endorsement
American College of Sports Medicine (ACSM)	X	X
American Cancer Society (ACS) <sup>a,b</sup>	X	X
American Academy of Physical Medicine and Rehabilitation (AAPMR) <sup>a,b</sup> and Foundation for PM&R	X	X
American Physical Therapy Association (APTA) and Academy of Oncologic Physical Therapy of the APTA <sup>a,b</sup>	X	X
American Congress of Rehabilitation Medicine		
American College of Lifestyle Medicine <sup>b</sup>		X
Canadian Society for Exercise Physiology (CSEP) <sup>a,b</sup>	X	X
Centers for Disease Control (CDC) <sup>b</sup>		X
Commission on Accreditation of Rehabilitation Facilities <sup>b</sup>		X
Exercise and Sports Science Australia (ESSA) <sup>a,b</sup>	X	X
German Union for Health Exercise and Exercise Therapy (DVGS) <sup>a,b</sup>	X	X
MacMillan <sup>a,b</sup>	X	X
National Cancer Institute (U.S.)		
National Comprehensive Cancer Network (NCCN) <sup>b</sup>		X
Royal Dutch Society for Physical Therapy (KNGF) <sup>a,b</sup>	X	X
Society for Behavioral Medicine (SBM) <sup>b</sup>		X
Sunflower Wellness <sup>a,b</sup>	X	X

<sup>a</sup>Roundtable Partner Organizations; <sup>b</sup>Organizations that provided official endorsement.

BOX 1. List of common acute, long-term, and late effects of cancer for review of evidence for therapeutic efficacy of exercise and subsequent exercise prescriptions

- Anxiety
- Bone health
- Cardiotoxicity
- Chemotherapy-induced peripheral neuropathy
- Cognitive function
- Depressive symptoms
- Falls
- Fatigue
- Health-related quality of life
- Lymphedema
- Nausea
- Pain
- Physical function
- Sexual function
- Sleep
- Treatment tolerance

expertise and were asked to contribute in one or more of the following areas: 1) role of exercise in cancer prevention and control; 2) efficacy of exercise to improve cancer-related health outcomes (acute, late, and long-term effects); and 3) translation of evidence into the clinical and community settings. The outcome of the work in each of these areas would be three separate, but related publications. The articles were circulated to professional organizations for review and to obtain their official endorsement.

This article will update evidence-based guidelines for exercise testing, prescription, and delivery in cancer survivors. As it was acknowledged that the general exercise recommendations put forward in 2010 may be unachievable for cancer survivors with physical limitations and that benefits may come from less exercise, a particular goal of the 2018 Roundtable was to develop more granular exercise prescriptions for distinct cancer-related health outcomes to better guide fitness and other health care professionals who train or care for cancer survivors. In the following sections we: 1) describe the evidence review process and decisions for generating exercise prescriptions for specific cancer-related health outcomes; 2) provide evidence-based prescriptions for frequency, intensity, time and type (FITT) for outcomes with sufficient evidence (as outlined below); and 3) provide updates to the 2010 guidelines around exercise testing and training, including special considerations and safety precautions, specific to cancer survivors. We

conclude by acknowledging the limitations of this latest Roundtable and suggest directions for future updates.

## UPDATE TO EVIDENCE-INFORMED EXERCISE PRESCRIPTIONS

### Overview

Two *a priori* decisions were made by consensus at the Roundtable meeting. The first was to develop a list of cancer-related health outcomes with a high degree of clinical relevance for which exercise may have therapeutic benefit (Box 1). The second was to focus the review of evidence primarily on traditional modalities of exercise, including aerobic, resistance, or combined aerobic plus resistance training on relevant health outcomes. A brief discussion on other modalities of exercise (e.g., yoga) is provided at the end of the article. Three additional decisions were made by the writing team early on in the writing process where it was agreed upon by consensus that: 1) components of physical fitness (e.g., aerobic capacity, muscular strength/endurance) would not be categorized as cancer health-related outcomes, but would be used to evaluate the adaptability and responsiveness of cancer survivors to specific modes of exercise training; 2) exercise prescriptions would only be generated for outcomes where there was sufficient evidence on the efficacy of exercise to improve a given outcome; 3) beyond each outcome, the exercise prescriptions could not be further specified by tumor type, phase of treatment or type of treatment due to the lack of sufficient evidence to do so in a robust manner. The implications of these limitations are discussed later in the article.

### Methodology for Evidence Review

To efficiently evaluate and provide a rich synthesis of the evidence, a review of published randomized controlled trials, systematic reviews, and meta-analyses for cancer-related health outcomes (Box 1) using Medline/PubMed, EMBASE, the Cochrane Central Register of Controlled Trials, CINAHL, the Physical Therapy Evidence Database (PEDro), and the Cochrane Collaboration. A search was conducted from June–August 2018, for articles published as of June 1, 2018, using standardized search terms for cancer and exercise, in




FIGURE 1—Decision tree on the level of evidence for effects (RCT) sufficient to merit an exercise prescription (FITT Rx).<sup>a</sup> Adapted with permission from Weggemans RM, Backx FJG, Borghouts L, et al. The 2017 Dutch Physical Activity Guidelines. *Int J Behav Nutr Phys Act.* 2018;15(1):58.

combination with search terms for the list of key cancer-related health outcomes (Supplemental Materials). For each outcome, two writing team members reviewed the resulting systematic reviews and meta-analyses to identify the most recent, relevant, and high-quality publications that could facilitate evaluation of the state of science around efficacy of exercise for a particular outcome. If no systematic reviews and meta-analyses were identified, the available randomized controlled trials (RCT) were reviewed.

A decision framework adapted from the Dutch Physical Activity Guidelines (12) was applied to first determine whether or not there was sufficiently strong evidence to conclude that exercise improved specific outcomes and in turn, warrant generation of an evidence-based FITT prescription (Fig. 1). Evidence for a given outcome was judged to be strong when there were a substantial number of RCT ( $\geq 5$ ), the aggregate sample size was large ( $n > 150$ ), and the beneficial effect of exercise was observed consistently across studies. For outcomes with a smaller number of RCT ( $< 5$ ) or where the overall effect of exercise was inconsistently observed or null, the level of evidence for an exercise benefit was judged as insufficient. For such cases, a FITT prescription was not generated, and only a summary of current evidence and research recommendations was provided.

During the review process, authors felt that an intermediate category should be considered for outcomes with  $> 5$  RCT and an aggregate sample  $> 150$ , yet heterogeneity for evidence of an effect in cancer survivors. In these cases, if there were a consistent number of high-quality RCT (designed and powered on the outcome of interest as the primary endpoint) and that reported findings that were congruent with established evidence on the same outcome in noncancer populations, the evidence of exercise benefit for these outcomes was judged as moderate and a FITT prescription was generated. However, more research would be needed to confirm the exercise recommendation for these outcomes in cancer survivors.

The next step was to define exercise prescriptions that conformed to the FITT formula and outlined the type and dose of exercise expected to improve a given outcome. To streamline the process, the evidence used to inform each exercise prescription was derived from high-quality systematic reviews and meta-analyses which also often provided separate analyses on efficacy by FITT components to identify the most optimal exercise type, time, intensity or frequency to improve a given outcome. However, if the quality of systematic reviews was deemed low (AMSTAR rating  $< 7$ ), individual high-quality RCT were used to derive the prescription. The FITT prescriptions and accompanying summary of the literature, which informed them, were presented to the full writing team during a series of conference calls for discussion and expert consensus (September to October 2018).

## Strong Evidence

For the following cancer-related health outcomes, there was consensus that benefit has been consistently demonstrated, and a FITT prescription was developed. Details of each FITT prescription, including any unique considerations, by outcome

are provided in Table 2. When interpreting and applying a prescription, the exercise professional should be mindful that the evidence is often disproportionately from trials in a single cancer type (i.e., breast), but the prescription is assumed to generalize across cancer types unless otherwise specified. In addition, studies often did not specifically target enrollment of individuals with the poorest initial values of an outcome (e.g., high fatigue, low bone density) thus the efficacy of a FITT prescription may or may not generalize to cancer survivors in greatest need.

**Anxiety.** A dose of moderate-intensity aerobic training three times per week for 12 wk or twice weekly combined aerobic plus resistance training for 6 to 12 wk can significantly reduce anxiety in cancer survivors during and after treatment (13–17). Based on sufficient evidence, it does not appear that resistance training alone reduces anxiety. There is not sufficient evidence to determine whether or not there is a dose response relationship between exercise intensity and changes in anxiety. Improvements in anxiety appear to be greater in supervised training programs or those having a larger supervised component than those that are predominantly unsupervised or home-based.

**Depressive symptoms.** Moderate-intensity aerobic training performed three times per week and for at least 12 wk or twice weekly combined aerobic plus resistance training lasting 6 to 12 wk can significantly reduce depressive symptoms in cancer survivors during and after treatment (18,19). Based on sufficient evidence, resistance training alone does not seem to be effective for this outcome. Based on results from high-quality trials of aerobic training, there may be a dose-response effect where higher volumes of aerobic exercise (90 min vs up to 180  $\text{min}\cdot\text{wk}^{-1}$ ) leads to better symptom reduction. Improvements in depressive symptoms appear to be greater in supervised training programs or those having a larger supervised component than those that are predominantly unsupervised or home-based.

**Fatigue.** For training programs that last at least 12 wk, engaging in moderate-intensity aerobic training three times per week can significantly reduce cancer-related fatigue both during and after treatment (20–23). Moderate-intensity combined aerobic plus resistance training sessions performed two to three times per week or twice weekly moderate-intensity resistance training may also be effective (21,24–26), and the latter particularly in prostate cancer (27). The effect of exercise was strongest for moderate- to vigorous-intensity exercise, whereas the effect for low-intensity training was weak, and this level of exercise is unlikely to reduce fatigue (22,24,26,28). Whether or not more exercise translates to less cancer-related fatigue remains unclear, though there is suggestive evidence that the reductions in fatigue are greater with exercise sessions longer than 30 min and programs longer than 12 wk compared with less exercise (29). There is insufficient evidence for a linear dose response since going beyond 150  $\text{min}\cdot\text{wk}^{-1}$  of aerobic exercise does not appear to result in the greatest reductions in fatigue. The efficacy of exercise for this outcome appears to be independent of the level of supervision and/or setting for training (21,27,30).

TABLE 2. Cancer-related health outcome with sufficient evidence for development of FITT prescription.

Outcome	Type <sup>a</sup>	Intensity	Duration (Min) or Sets (Reps)	Frequency (Sessions per Week)	Length (Weeks)	Setting (Supervised, Home-Based or Combination)	Dose Response	Special Considerations	Evidence Primarily from These Cancer Types.
Anxiety	Aerobic	60%–80% HRmax 60%–80% VO <sub>2</sub> max RPE 13–15	30–60	3	12	Supervised more effective	Moderate to vigorous may be more effective than light to moderate	Not known	Breast (majority), prostate, colorectal, gynecological (ovarian, endometrial, cervical), head and neck, lung, hematological cancer
Depressive Symptoms	Resistance	Efficacy not demonstrated	NA	NA	NA	NA	NA	NA	
	Aerobic + Resistance	60%–80% HRmax 60%–80% VO <sub>2</sub> max RPE 13–15	20–40	2–3	6–12	Supervised or combination of supervised & home-based	None observed	Not known	
	Aerobic	65%–85% 1-RM	2 sets 8–12 reps 30–60	2–3	6–12	Supervised more effective	Benefit up to 180 min/wk	Not known	Breast (majority), prostate, colorectal, hematological
Fatigue	Resistance	Efficacy not demonstrated	NA	NA	NA	NA	NA	NA	
	Aerobic + Resistance	60%–80% HRmax 60%–80% VO <sub>2</sub> max RPE 13–15	20–40	3	12	Supervised or combination of supervised & home-based	None observed	Not known	
	Aerobic	65%–85% 1-RM	2 sets 8–12 reps 30	2–3	6–12	Supervised and unsupervised appear similarly effective	No dose response by intensity; possible > benefits with ↑ duration & length of program	No evidence of benefits from light intensity	Breast (majority), prostate, mixed
Health-related Quality of Life	Resistance	60% 1-RM RPE 12	2 sets 12–15 reps 30	2	12	Supervised and unsupervised appear similarly effective	None observed	Not known	
	Aerobic + Resistance	65% HRmax 45% VO <sub>2</sub> max RPE 12	2 sets 12–15 reps 30	3	12	Supervised and unsupervised appear similarly effective	None observed	Not known	
	Aerobic	60%–80% HRmax RPE 11–13	2 sets 12–15 reps 30	2–3	12	Supervised more effective	None observed	NA	Breast (majority), prostate, colorectal, lung, head and neck, bladder, gynecological, mixed, hematological cancer, hematopoietic stem cell transplant
Lymphedema <sup>b</sup>	Resistance	60%–75% 1-RM RPE 13–15	2–3 sets 8–15 reps 20–30	2–3	12	Supervised or combination of supervised & home-based	None observed	NA	
	Aerobic + Resistance	60%–80% HRmax RPE 11–13	2 sets 8–15 reps NA	2–3	12	Supervised more effective	None observed	AT and RT combined most effective	
	Aerobic	60%–80% 1-RM RPE 12–14 NA	2 sets 8–15 reps NA	2–3	12	Supervised more effective	None observed	Generally safe, as no significant increase in number of lymphedema-related adverse events reported in RCT of aerobic exercise	Breast cancer-related lymphedema only
	Resistance	60%–70% 1-RM RPE 15	1–3 sets 8–15 reps	2–3	52	All of the interventions reviewed started with supervision to teach the exercises.	NA	Start resistance a low weight and progress slowly	
	Aerobic + Resistance	NA	NA	NA	NA	NA	NA	NA	



TABLE 2. (Continued)

Outcome	Type <sup>a</sup>	Intensity	Duration (Min) or Sets (Reps)	Frequency (Sessions per Week)	Length (Weeks)	Setting (Supervised, Home-Based or Combination)	Dose Response	Special Considerations	Evidence Primarily from These Cancer Types.
Physical Function	Aerobic	60%-85% HRmax 60%-85% VO <sub>2max</sub> RPE 12-13	30-60	3	8-12	Supervised more effective	If unsupervised requires higher weekly exercise expenditure (unclear what the threshold is)	NA	Breast (majority), prostate, colorectal, lung, bladder, head and neck, hematological cancer, hematopoietic stem cell transplant
	Resistance	60%-75% 1-RM RPE 13-15	2 sets 8-12 reps	2-3	8-12	Limited evidence to determine benefit of unsupervised	NA	NA	
	Aerobic + Resistance	60%-85% HRmax 60%-85% VO <sub>2max</sub> RPE 12-13	20-40	3	8-12	Both supervised & home-base suitable in older adults	NA	Community-based interventions that met in groups & used behavior change strategies may produce larger effects in older adults	

<sup>a</sup>Mode: For aerobic activity, this includes walking, cycling ergometer and other forms of traditional aerobic exercise; for resistance, this includes machine-based and free weights.

<sup>b</sup>Recommendation for breast cancer-related lymphedema only. Abbreviations: NA, not applicable.

**Health-related quality of life.** Combined moderate-intensity aerobic and resistance exercise performed two to three times per week for at least 12 wk results in improvements in health-related quality of life (HRQoL) both during and after treatment (17,31,32). The benefit of combined aerobic plus resistance training programs appears more potent than programs consisting of only aerobic or resistance training (32). Though enough evidence favored the efficacy of exercise to improve HRQoL, it should be noted that this particular outcome is a construct that encompasses many factors and may have a more variable response than individual domains. For example, improvements seem to be more robust if the physical functioning domain of HRQoL is the primary outcome, and this will be covered separately. Improvements in HRQoL appear to be greater in supervised training programs or those having a larger supervised component than those that are predominantly unsupervised or home-based.

**Lymphedema.** There is a history of clinical recommendations to refrain from aerobic or resistance training to avoid onset or exacerbation of lymphedema (33,34). For this particular outcome, an evidence-based exercise prescription is designed for safety or no harm versus the benefit of exercise to prevent lymphedema or improve lymphedema symptoms and limited to addressing upper-extremity breast cancer-related lymphedema (35-37). For resistance training, a general progressive program focused on large muscle groups performed two to three times per week, with the principle of “start low, progress slow” is safe when supervised by a fitness professional (38-42). Insufficient evidence exists to conclude whether or not starting a resistance training program *without* supervised instruction is safe for women with or at risk for lymphedema after breast cancer. To date, there is insufficient evidence to draw conclusions for aerobic exercise. In general, aerobic exercise seems to be safe, with no significant increase in number of lymphedema-related adverse events reported in studies investigating aerobic exercise (17). Based on preliminary evidence (43), the effects seen in breast cancer may not translate for lymphedema following head and neck, bladder, melanoma, gynecologic, or other cancer sites.

**Physical function.** Moderate-intensity aerobic training, resistance training or combined aerobic plus resistance training performed three times weekly for 8 to 12 wk can significantly improve self-reported physical function (31,32,44). Broadly, supervised exercise appears to be more effective than unsupervised or home-based interventions (32), although unsupervised programs may be effective in older cancer survivors (44). Further, there is some evidence to suggest that if the intervention is unsupervised, physical function could improve with higher weekly energy expenditure (MET·h·wk<sup>-1</sup>) but not weekly exercise duration (min·wk<sup>-1</sup>) (32). It should be noted that these results are based on self-reported physical function, not objective measures where the evidence base on these outcomes remains immature and more challenging to aggregate due to the variation and limitations of assessment techniques.

## Moderate Evidence

**Bone health.** Two recent systematic reviews in cancer survivors concluded that across all trials the evidence for exercise to improve bone health is inconsistent (45,46), though RCT that were designed with bone health as the primary outcome were largely consistent with the exercise recommendations in the ACSM Position Stand for exercise and bone health (47). In cancer, the majority of evidence is derived from trials in breast and prostate cancer patients in the postadjuvant treatment setting, which indicates that a 1-yr supervised program of combined moderate-vigorous intensity resistance *plus* high-impact training (i.e., exercise that generates ground reaction forces above three to four times body weight) performed 2 to 3 d·wk<sup>-1</sup> is the most consistently efficacious modality of exercise to improve bone health (e.g., slow loss or slightly improve bone mineral density at the lumbar spine and hip). In contrast, aerobic training, particularly walking, does not appear to provide a sufficient stimulus to improve bone outcomes, results that are consistent with RCT in persons without cancer. There is insufficient evidence to determine if resistance training alone improves bone outcomes.

In noncancer populations, whether or not resistance plus impact training programs are safe for individuals with osteoporosis remains controversial (48), thus, at this time, this exercise prescription may not be safe for cancer survivors with bone fragility associated with osteoporosis or bony metastases in the hip or spine (see Exercise Safety and Training Tolerance). Furthermore, it may not be appropriate for individuals with joint/orthopedic issues and/or stability problems who may be better served by an exercise program aimed at reducing fall risk. Further research is needed to confirm that this recommendation is effective and safe for cancer survivors.

**Sleep.** Two recent systematic reviews in cancer survivors (49,50) provided mixed evidence for overall sleep quality, indicating either a positive effect of walking (49) or no effect of exercise (50). Four recent RCT not included in either systematic review have shown consistent evidence of small to moderate effect sizes on overall sleep quality for aerobic training (51), in addition to walking specifically (52,53) and one study reported evidence of benefit for resistance training (31). In noncancer populations, there is strong evidence that moderate to vigorous intensity aerobic training is associated with better overall sleep quality in the general population (54), and there was some evidence showing benefit for specific characteristics of sleep, such as total sleep time, sleep onset latency, and sleep efficiency. Overall, moderate-intensity aerobic training, particularly walking, three to four times per week, for 30 to 40 min per session over 12 wk is recommended.

## Insufficient Evidence

Insufficient evidence for a specific outcome does not mean that cancer survivors facing these cancer-related health outcomes will not benefit in other ways from engaging in physical activity or should remain sedentary. Rather, there

is insufficient evidence showing that exercise is beneficial for these specific outcomes based on current evidence, and this creates an obvious gap in knowledge the research community must fill.

**Cardiotoxicity.** The ability of exercise to prevent or ameliorate cardiotoxicity is an emerging field of research and promising results for a protective effect of exercise in animal models and some novel evidence in humans for cardiac function (55), including measures of left ventricular function (55), and vascular endothelial function, measured as flow-mediated dilation (56). More research is needed to understand the impact of various types of cancer, cancer treatments, and exercise prescriptions on both cardiac and vascular function.

**Chemotherapy-induced peripheral neuropathy.** To date, there are too few high-quality trials to interpret the potential of the benefits of exercise for preventing and/or managing chemotherapy-induced peripheral neuropathy and related side effects, such as balance impairment and falls (57). In general, exercise appeared safe in a few published studies; however, the degree of improvement varied for this outcome, and other related outcomes, such as mobility and balance. Future research should include well-controlled exercise interventions with chemotherapy-induced peripheral neuropathy as a primary outcome, using both patient-reported outcomes and objective assessment of neuropathy, balance, and mobility, and also rigorously evaluate the safety of training in this group due to the known risk of falls.

**Cognitive function.** Although promising results from animal studies are emerging for a protective effect of aerobic training on cancer treatment-related changes in cognitive function, to date, the evidence in humans is limited (58). The majority of human studies, so far, have been conducted in breast cancer survivors, using self-reported measures of cognitive function, and report inconsistent results. Although there is compelling evidence of a positive effect of exercise on cognitive function in older adults and other clinical populations (59), specific to cancer, more research is needed in cancer survivors where cognitive function is the primary outcome, and this outcome is assessed by both self-reported and standardized objective measures of cognitive function.

**Falls.** There are no randomized controlled trials to date in cancer survivors with falls as a primary endpoint. There are several challenges to this type of research, including the relatively rare occurrence of falls and the large sample and time needed to observe a change in falls from an intervention. Similarly, the causes of falls associated with cancer treatment have not been fully characterized and may be due to more than an acceleration of the risks associated with age-related falls (i.e., muscle weakness and poor balance), but may also be due to treatment-related toxicities, such as hearing loss, ataxia, peripheral neuropathies, and fatigue, creating a challenge to develop new exercise-based approaches to fall prevention. In the absence of any evidence-based fall prevention studies in cancer survivors, it seems reasonable to consider standard fall prevention exercise approaches that reduce the risk of age-related falls for cancer

survivors with a fall history to at least reduce the risk of falls that may be associated with advanced age (60,61).

**Nausea.** Although reduction in nausea is a commonly reported benefit of exercise during chemotherapy, there is limited data from high-quality trials with nausea as a primary endpoint to support this finding outside of an early study demonstrating a reduction by Winningham et al. (62). Future research should examine the effect of well-controlled exercise interventions on nausea in highly emetogenic treatment regimens to determine if there is an effect. These studies should account for antiemetic use, duration and intensity of nausea, and function.

**Pain.** To date most published controlled trials in cancer survivors have examined nonspecific pain, and included pain as a secondary outcome, which limits the interpretation of the research. There is early evidence from two high-quality controlled trials where pain was the primary outcome that a combined home-based aerobic plus supervised resistance training intervention in women with breast cancer significantly reduced arthralgia associated with aromatase inhibitor therapy (63) and a supervised resistance training intervention focused on the upper extremity significantly reduced in shoulder pain in individuals with head and neck cancer (64). However, more research is needed specifically focused on cancer-related pain. Survivors may be able to exercise with pain that is tolerable and not worsened by exercise; however, modification or omission of individual exercises that exacerbate pain may be necessary.

**Sexual function.** The majority of research to date has been in men with prostate cancer (65,66), and there is insufficient evidence from controlled trials investigating the effect of exercise on sexual function during or after cancer therapy. Early promising results for a positive effect of exercise on sexual function among prostate cancer patients treated with androgen deprivation therapy (ADT) has been reported in some trials (67,68) but not others (69). Although there is compelling evidence of a positive effect of exercise on sexual function in the general population, for both women (70) and men (71), the significant effects of cancer therapies on hormonal and anatomical/functional changes (i.e., nerve sparing vs non-nerve-sparing surgeries in prostate cancer) preclude the ability to extrapolate those findings to the cancer population.

**Treatment tolerance.** Treatment tolerance (i.e., completion of or adherence to planned therapy) is a complex outcome that likely varies by cancer type, treatment modality (i.e., radiation therapy, chemotherapy, hormone therapy, immunotherapy), and even specific drug(s) and protocols. Consequently, determining the effects of exercise on “treatment tolerance” is a challenging goal and may make it difficult to achieve a high degree of generalizability. A recent systematic review of exercise and chemotherapy completion rate concluded that, although promising, the evidence for an exercise benefit to chemotherapy tolerance is insufficient (72). The effects of exercise on treatment tolerance for radiation therapy, hormonal therapy (i.e., aromatase inhibitors or ADT), targeted therapies, or immunotherapy are currently unknown. To understand the effect of exercise on treatment tolerance,

studies would need to be conducted for each combination of cancer type and treatment regimen.

## Limitations

The FITT prescriptions provided for several outcomes aim to serve as guidelines for fitness and health care professionals working with cancer survivors. However, the development of these guidelines had some key limitations that should be kept in mind when working with individual clients. The majority of available literature is still in the most common cancers, namely, early stage breast cancer and prostate cancers, which limits the ability to extrapolate their findings to other cancer types or advanced cancers. Though, it could be reasonably assumed that in the absence of any unique safety concerns for survivors of other types of cancer that the efficacy of exercise on various outcomes would be similar for survivors with early stage cancers other than breast and prostate cancer. However, differences among cancer survivors by cancer type are known to exist (i.e., demographics, prognosis, treatments received, and associated side effects) and further, there is very little information regarding the feasibility, safety or benefits of exercise in individuals living with advanced cancer. More research is needed to advance the level of specificity available in the exercise oncology literature to serve a broader range of cancer types and stages (73). In addition, there has historically been incomplete reporting of compliance to a prescribed FITT program in published studies, likely because tracking and quantifying compliance can be burdensome (74–76). Thus, it remains possible that some proportion of cancer survivors may not be able to tolerate the evidence-based FITT, as explicit reporting of adjustments to the exercise prescription (i.e., dose modification) based on tolerance of individuals has been limited. Thus, the fitness or health care professional should monitor for early signs of poor tolerance to training and adjust the dose of exercise accordingly even if this means dropping below recommended training volumes. Furthermore, understanding the specific efficacy of exercise for a particular outcome is hampered by the fact that the majority of research to date does not consider the training principle of initial values by limiting enrollment to individuals experiencing the specific outcome of interest (i.e., those with sleep issues or high level of fatigue), rarely examines potential moderators of the exercise response (i.e., baseline functional capacity), and often draws conclusions about outcomes that are secondary to the intended design of the study. Finally, a further understanding of dose response is limited by a paucity of trials that directly compare two or more levels of exercise training (e.g., high-intensity vs low-intensity exercise) on cancer-related outcomes (77) or compare settings (e.g., supervised vs home-based) (78). Clearly, more research to address these knowledge gaps is warranted so that recommendations can continue to improve in scope and specificity.

## Other Exercise Modes

There is also increasing interest in the safety and efficacy of types of exercise that fall outside of traditional modes of



aerobic and resistance training. In a recent systematic review of the role of yoga in symptoms management for cancer survivors, yoga both during and after cancer treatment was reported to improve quality of life and fatigue, whereas further research is needed to confirm the observed potential to improve sleep, depressive symptoms, anxiety/distress, and cancer-related cognitive change (79). However, the literature is reflective of practices for many different types of yoga, including those that incorporate nonexercise features, such as breathing or meditation, and which use a wide variety of prescriptions (e.g., frequency, duration, with/without home practice), making it difficult to generate a definitive prescription. Although there is also insufficient evidence at this time for a definitive prescription around safety and efficacy for other types of exercise for cancer survivors, such as dragon boating (80,81), recreational sports (82), wall/rock climbing (83), triathlon (84), or high-intensity interval training (85), research is on-going.

## EFFECTS OF CANCER TREATMENT AND ADVERSE EFFECTS RELEVANT TO EXERCISE

To best evaluate a cancer survivor's exercise tolerance and prescribe a safe and effective exercise program, it is necessary for fitness professionals to know about the type and extent (i.e., stage) of cancer a person has. Fitness professionals must also be familiar with the common treatment approaches to cancer, the side effects and symptoms these treatments can cause, and the subsequent impact on exercise tolerance (Table 3). The treatment approach used will differ by type of cancer, stage of disease, cancer subtype, patient health, and many other considerations. Treatment modalities may include a

combination of surgery, radiation, and systemic therapies, including cytotoxic chemotherapy, newer targeted agents including immune checkpoint inhibitors, and hormonal therapy. When individuals are on active cancer treatment, working closely with the oncology treatment team is recommended, as treatment approaches change frequently and understanding the side effects of newer treatments continues to evolve.

The impact of cancer treatment on exercise tolerance may further depend upon the prediagnosis health and functional capacity of the individual. Furthermore, fitness professionals should be aware of, and respectful of, the fact that individuals diagnosed with cancer commonly have many concerns, such as life expectancy, employment issues, and family matters, that may limit prioritization of exercise in their lives.

## MEDICAL CLEARANCE AND EXERCISE TESTING

Both the diagnosis of cancer and curative cancer treatments may affect the underlying safety of exercise training. Guidance for the indications of medical clearance before exercise testing and/or training, as well as how exercise testing should be adapted for cancer survivors, can be useful for creating a safe and effective exercise prescription. Where this information is available elsewhere and to avoid redundancy, the reader will be referred to specific publications.

### Medical Clearance Before Exercise

Given the diversity of tumor types and side effect of different cancer treatments, including the potential acceleration of

TABLE 3. Potential impact of cancer treatments on exercise tolerance and safety.

	Surgery	Chemotherapy	Radiation	Anti-Hormonal Therapy (Surgical or Pharmaceutical)	Targeted Therapy or Immunotherapy <sup>a</sup>
Cardiovascular changes	Cardiac damage or increased CVD risk	√	√	√	√
Endocrine changes					
	Worsening bone health	√	√	√	
	Changes in body composition (weight gain)	√		√	
	Changes in body composition (weight loss/muscle mass loss)	√	√	√	√
Gastrointestinal changes					
	Nausea	√			√
	Diarrhea	√	√		√
	Altered GI function	√	√		√
Immune changes	Impaired immune function and/or anemia	√	√	√	√
Metabolic changes					
	Development/worsening of metabolic syndrome	√		√	√ <sup>b</sup>
Neurological changes					
	Peripheral Neuropathy	√			
	Cognitive changes	√ (brain surgery)	√	√	
Pulmonary changes	Altered lung function or pneumonitis	√ (lung surgery)	√		
Skin changes					
	Redness, irritation		√		
	Rashes		√		√
	Reduced ROM	√ (by healing at surgical site)	√		
Fatigue		√	√	√	√
Lymphedema <sup>c</sup>		√	√		
Pain					
	General	√	√	√	√
	Myalgia/arthralgia		√	√	√

Adapted from Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc.* 2010;42(7):1409–26.

<sup>a</sup>Depends on type or target of agent.

<sup>b</sup>Especially common with PI3kinase inhibitors.

<sup>c</sup>Can occur in any type of cancer when and where lymph nodes are surgically resected and/or radiation over lymph nodes.

CVD, cardiovascular disease; GI, Gastrointestinal.

TABLE 4. Adapted national comprehensive cancer network triage approach based on risk of exercise-induced adverse events.

Description of Patients No comorbidities	Evaluation, prescription, and programming recommendations No further preexercise medical evaluation <sup>a</sup> Follow general exercise recommendations
Peripheral neuropathy, arthritis/musculoskeletal issues, poor bone health (e.g., osteopenia or osteoporosis), lymphedema	Recommend preexercise medical evaluation <sup>a</sup> Modify general exercise recommendations based on assessments Consider referral to trained personnel <sup>b</sup>
Lung or abdominal surgery, ostomy, cardiopulmonary disease, ataxia, extreme fatigue, severe nutritional deficiencies, worsening/changing physical condition (i.e., lymphedema exacerbation), bone metastases	Preexercise medical evaluation <sup>a</sup> and clearance by physician before exercise Referral to trained personnel <sup>b</sup>

<sup>a</sup>Medical evaluation—per NCCN guidelines for specific symptoms and side effects. <sup>b</sup>Rehabilitation specialists (i.e., physical therapists, occupational therapists, physiatrists) and certified exercise physiologists (i.e., ACSM Certified Clinical Exercise Physiologist (ACSM-CEP), Canadian Society for Exercise Physiology Certified Exercise Physiologist (CSEP-CEP), Exercise & Sport Science Australia Accredited Exercise Physiologist (ESSA-AEP)).

cardiovascular disease (CVD), the question of whether or not cancer survivors require medical clearance (i.e., approval from a medical professional to engage in exercise) before starting an exercise program is always relevant. Recently, the ACSM updated its preparticipation exercise guidelines for all persons in an attempt to reduce barriers to exercise by removing a requirement for medical clearance for individuals whose risk of an adverse cardiac event during exercise are low, including exercise naïve persons (86). Preparticipation guidelines for evaluating the need for medical clearance for noncancer comorbidities should be applied in cancer survivors to minimize risks of adverse exercise-related events. The ACSM preparticipation guidelines do not explicitly address risks for adverse events and/or injury during exercise that are specific to the adverse effects of cancer treatment. Therefore, we have referred to the National Comprehensive Cancer Network (NCCN) Survivorship Guidelines (8) to frame recommendations for when medical clearance and/or further medical

evaluation by a medical professional is indicated, as well as the level of supervision during exercise training for cancer survivors to ensure safety based on the disease- and treatment-related side effects (Table 4).

### Exercise Testing

Ideally, cancer survivors should receive a comprehensive assessment of all components of health-related physical fitness (i.e., cardiorespiratory fitness, muscle strength and endurance, body composition, and flexibility), with some specific cancer-specific considerations (Box 2), to individualize an exercise prescription. However, requiring a comprehensive physical fitness assessment before starting exercise may create an unnecessary barrier to starting activity. For this reason, no assessments are required to start low-intensity aerobic training (i.e., walking or cycling), resistance training with gradual progression, or a flexibility program in most survivors. Medical clearance may still be indicated as previously described depending on exercise and health history and presence of cardiovascular, renal, or metabolic symptoms (86).

#### BOX 2. Exercise testing recommendations:

- Standard exercise testing methods are generally appropriate for patients with cancer who do not require preexercise medical evaluation or who have been medically cleared for exercise with the following considerations:
- Be aware of a survivor's health history, comorbid chronic diseases, and health conditions, and any general exercise contraindications before commencing health-related fitness assessments or designing the exercise prescription (86).
- Be familiar with the most common toxicities associated with cancer treatments including increased risk for fractures and cardiovascular events, along with neuropathies or musculoskeletal morbidities related to specific types of treatment
- Health-related fitness assessments may be valuable for evaluating the degree to which components of fitness have been affected by cancer-related fatigue or other commonly experienced symptoms that impact function (87)
- In principle, there is no evidence that the level of medical supervision required for symptom-limited or maximal cardiopulmonary exercise testing need to be different for patients with cancer than for other populations (86).
- The evidence-based literature indicates 1-repetition maximum (RM) testing is safe among survivors of breast and prostate cancer without bony metastases (6)
- Among patients with bony metastases or known or suspected osteoporosis routine assessments of muscle strength and/or endurance involving musculature that attaches to and/or acts on a skeletal site that contains bone lesions should be avoided (88). For example, 1-RM testing for leg strength (e.g., leg press) should be avoided in patients who have bony metastases in the proximal femur (i.e., hip) or vertebrae. Other sites where lesions are absent could be tested. In this example, if the patient had no lesions in the upper body, 1-RM for a chest press or 1-RM for a seated row might be feasible, given no other contraindications. Medical clearance from a physician (i.e., orthopedic or radio-oncology) may be mandatory depending on scope of practice or protocols at a specific site/facility.
- Older survivors and/or survivors treated with neurotoxic chemotherapy (typical for breast, colon, lung, ovarian cancers) may especially benefit from a standard assessment of balance and mobility to assess fall risk (89)
- CVD has become a competing cause of morbidity and mortality for survivors of cancer with a favorable prognosis (90). Given the potential for underlying CVD, cancer survivors should be screened for evident or underlying CVD using the ACSM preparticipation guidelines (see below) and if implicated have a cardiopulmonary exercise test before beginning an exercise program (91).

### EXERCISE SAFETY AND TRAINING TOLERANCE

**Safety of exercise training.** The overall conclusion from the 2010 Roundtable was that exercise is generally safe for cancer survivors (6) and this has not changed based on the majority of studies conducted since that time. It should be recognized that the majority of available evidence on the safety and efficacy of exercise during and after cancer treatment is derived from RCT of supervised and/or home-based prescribed exercise, and trials in breast cancer survivors (17,31,32). Hence, the individuals enrolled in studies commonly meet prespecified eligibility criteria for age, comorbidities, physical ability, largely based out of academic and/or medical centers, and were willing to participate in research. This often results in a sample that is healthier or with higher physical function and exercise motivation that may not fully generalize to the broader population of cancer survivors. Depending on the nature and extent of a survivor's presenting problems, they may not be able to adequately and/or safely engage in the levels of exercise outlined in this recommendation. In these cases, we again refer to the NCCN guidelines (Table 4). Physical therapy or medical evaluation might be a bridge to inform appropriate modifications to an individual's exercise program and/or correct toxicities, impairments,

and limitations that prevent a survivor from working toward recommended levels of exercise.

**Exercise tolerance.** Exercise has well-established health benefits in persons without cancer, thus, a key consideration in exercise trials has been whether or not cancer survivors can tolerate the doses of exercise known or hypothesized to effectively improve physical fitness and in turn, associated cancer-related outcomes. Research, to date, supports the potential of cancer survivors to respond positively to an exercise training stimulus by improving individual components of physical fitness, including cardiorespiratory fitness (i.e.,  $\dot{V}O_{2peak}$ ) (92), muscular strength and endurance (93,94), and body composition (95). However, an individual's response to a given exercise stimulus may vary due to the direct effects of cancer treatments on physiological systems (i.e., anemia), side effects of cancer treatment (i.e., cancer-related fatigue may lower exercise tolerance), or demographics factors (i.e., age) (96). Furthermore, during active treatment an individual's ability to tolerate exercise may fluctuate from day to day or week to week. Understanding of these interactions is a topic of ongoing research, especially with the emergence of novel therapies.

Specific to cardiorespiratory fitness, during chemotherapy treatment, there is a well-documented decline in cardiorespiratory fitness, as measured by  $\dot{V}O_{2peak}$  or 6-min walk test (4). Randomized trials of aerobic exercise during adjuvant chemotherapy demonstrate a preservation of, or an improvement in, cardiorespiratory fitness, especially in those with low initial values (92), whereas others report better improvement in those with higher initial values (97). Aerobic exercise during adjuvant chemotherapy does not appear to stimulate greater production of red blood cells (98), so improvements in cardiorespiratory fitness are contingent on other central (i.e., cardiac function, plasma volume) and peripheral adaptations (i.e., improved vascularization and mitochondrial enzyme function) (99).

Specific to muscular strength, loss of muscle strength and endurance is common due to deconditioning or as a side effect of cancer treatment. For example, ADT, which is commonly used as treatment for prostate cancer, results in an abrupt loss of lean body mass accompanied by a reduction in muscle strength and endurance (100,101). In the absence of the anabolic drive from testosterone, men on ADT may not be able to build lean mass in response to resistance training; however, several trials that have employed resistance training, or combined with aerobic training, have reported small, but statistically significant improvements in lean body mass after 12 to 36 wk of training (102). Although sarcopenia is related to muscle weakness and contributes to poor functioning in older adults, neuromuscular contributions explain up to 50% of variation in muscle strength in older adults, thus resistance training in the setting of ADT or deconditioning may still effectively improve muscle strength in the absence of gains in muscle mass (103).

Specific to body composition, maintenance of body weight can be difficult during treatment for some cancers, where loss

of weight and lean body mass is a common concern, such as advanced colon, lung cancer, and pancreatic cancer (104), whereas weight gain can be a common side effect of chemotherapy and antiestrogen therapy for breast cancer or antiandrogen therapy for prostate cancer (105). Moreover, obesity is a risk factor for multiple cancers, including postmenopausal breast, renal cell, and endometrial cancer, thus these survivors are more likely to be overweight or obese at the time of diagnosis (106). In cases where weight and lean body mass loss may be a side effect of treatment, the fitness professional needs to ensure that exercise training is not creating an excess energy deficit (i.e., energy expenditure exceeds adequate dietary energy and nutrient intake) that contributes to weight loss and can aggravate fatigue (107,108). Working with a trained oncology dietician who can advise on dietary modifications that would support adequate fuel availability and replacement during and after exercise, respectively, may be prudent. For cases where survivors may be prone to weight gain and/or obesity, the exercise professional should be aware of the safety considerations related to exercise, including orthopedic limitations and cardiovascular disease risk (86,109). If weight loss is implicated in the health goals for these individuals, it may be prudent for the exercise professional and/or survivor to partner with a registered dietician to provide dietary recommendations that can complement an exercise program.

Specific to musculoskeletal flexibility, surgery can result in temporary or more permanent reductions in joint range of motion, and extensibility of muscle, tendon, fascia, and skin. Exercise professionals should be aware of surgical sites and if abnormal movement patterns are observed, adapt the proposed movements to avoid placing abnormal strain on other body structures and consider referral to physical therapy in efforts to address restrictions.

## IMPLEMENTING FITT PRESCRIPTIONS IN PRACTICE

Based on the current literature, an effective exercise prescription that most consistently addresses health-related outcomes experienced due to a cancer diagnosis and cancer treatment includes moderate-intensity aerobic training at least three times per week, for at least 30 min, for at least 8 to 12 wk. The addition of resistance training to aerobic training, at least two times per week, using at least two sets of 8 to 15 repetitions at least 60% of one repetition maximum, appears to result in similar benefits (Box 3). Exercise programs that only prescribe resistance training are also efficacious at improving

BOX 3. Expected patient benefits from exercise training by mode

Aerobic	Resistance	Aerobic plus Resistance
Reduced anxiety	Less fatigue	Reduced anxiety
Fewer depressive symptoms	Better QoL	Fewer depressive symptoms
Less fatigue	No risk of exacerbating lymphedema	Less fatigue
Better QoL		Better QoL
Improved perceived physical function	Improved perceived physical function	Improved perceived physical function

most health-related outcomes, though for some specific outcomes the evidence is either insufficient or suggestive that resistance training alone may not be enough (e.g., depressive symptoms). Exercise programs that were supervised appear to be more effective than strictly unsupervised or home-based programs, though it is unclear whether or not this is because a higher dose of exercise may be better achieved with supervised training or from other attributes of this setting (i.e., more attention, motivation, reinforcement, selection bias). Although a variety of professionals delivered supervised interventions in the research literature (e.g., exercise physiologists, certified exercise instructors, nurses, physical therapists) determining the type of professional that could maximize outcomes was beyond the scope of this article and the available evidence.

However, the fitness professional should be prepared to create an exercise program that meets their clients' needs. A customized program may not yet resemble or reach the exercise programs recommended in these guidelines, such that a goal may be to strive toward preparing the client to engage in recommended types and levels of exercise over their lifetime as outlined in the 2018 Physical Activity Guidelines for Americans (54). There is consistent observational evidence that engaging in physical activity after a cancer diagnosis reduces the risk of cancer-specific and all-cause mortality

for individuals diagnosed with early stage breast, colorectal, and prostate cancer (110). Special considerations and modifications to exercise programs have been adapted from the NCCN guidelines (Table 5). Finally, as part of the ACSM Roundtable efforts, oncologists are being asked to "Assess, Advise, and Refer," to connect cancer survivors to the most appropriate available exercise programming. A registry of programming is available at [www.exerciseismedicine.org/movethruca](http://www.exerciseismedicine.org/movethruca) (124).

## SUMMARY AND FUTURE DIRECTIONS

The 2018 ACSM Roundtable recommendations were made possible due to the increase in the availability of high-quality randomized controlled trials of exercise in cancer survivors published after the 2010 recommendations were issued. This allowed for the development of more specific evidence-based exercise prescription to improve common side effects of a cancer diagnosis and treatment, namely anxiety, depressive symptoms, fatigue, health-related quality of life, and physical function, along with safety of exercise training in persons with or at risk of breast cancer-related lymphedema. Future research is needed to determine the efficacy of exercise to improve other outcomes, including those identified here under the emerging or

TABLE 5. Exercise programming considerations for specific cancer survivors.

Considerations	Recommendations
Bone loss/bone metastases:	<ul style="list-style-type: none"> <li>• Avoid contraindicated movements that place an excessively high load on fragile skeletal sites. These include the following: high-impact loads, hyperflexion or hyperextension of the trunk, flexion or extension of the trunk with added resistance, and dynamic twisting motion</li> <li>• Specific guidance on how to modify exercise programs based on the site of bony lesions is provided elsewhere (88,111)</li> <li>• Preventing falls must also be a goal of therapy, since falls play an important role in fracture etiology (112).</li> <li>• Be aware of signs and symptoms of bone metastases in survivors, as well as common locations where these occur (i.e., spinal vertebrae, ribs, humerus, femur, pelvis). Bone pain can be an initial sign of skeletal metastases thus, exercise trainers should refer survivors who report pain back to the medical team for clinical evaluation before continuing exercise</li> </ul>
Lymphedema	<ul style="list-style-type: none"> <li>• To date, there is insufficient evidence to support or refute this clinical advice to wear a compression garment during exercise to prevent or reduce symptoms of breast cancer-related upper body lymphedema. Therefore, it is recommended that exercise professionals provide this information as part of client education and defer to an individual client's preference regarding use of a compression sleeve.</li> <li>• Being overweight or deconditioned have been associated with a higher risk of developing cancer-related lymphedema in observational studies, at this time there is insufficient evidence that weight loss or improving aerobic fitness can lower the risk of developing cancer-related lymphedema (113).</li> </ul>
Older adults	<ul style="list-style-type: none"> <li>• Physical problems reported by cancer survivors, such as cognitive difficulty, neuropathy, sarcopenia, muscle weakness, slowing, and fatigue, may be similar to those of older people without cancer, but cancer treatment can accelerate these declines (114–116)</li> <li>• Exercise professionals will need to combine ACSM guidelines on exercise programming for older adults (117) with the recommendations in this publication.</li> <li>• Integrate fitness and functional assessments before beginning an exercise program to more accurately determine baseline functional abilities.</li> </ul>
Ostomy	<ul style="list-style-type: none"> <li>• Empty ostomy bag before starting exercise</li> <li>• Weight lifting/resistance exercises should start with low resistance and progress slowly under the guidance of trained exercise professionals. People with an ostomy may be at an increased risk of parastomal hernia. To regulate intra-abdominal pressure, correct lifting technique and good form is required. Avoid use of a Valsalva maneuver (118,119).</li> <li>• Modify any core exercises which cause excessive intra-abdominal pressure, namely a feeling of pressure or observed bulging of the abdomen.</li> <li>• Those with an ileostomy are at increased risk of dehydration. Get medical advice on ways to maintain optimum hydration prior, during and after exercise.</li> <li>• Those doing contact sports or where there is a risk of a blow to the ostomy may wish to wear an ostomy protector/shield.</li> </ul>
Peripheral neuropathy	<ul style="list-style-type: none"> <li>• Stability, balance, and gait should be assessed before engaging in exercise; consider balance training as indicated</li> <li>• Consider alternative aerobic exercise (stationary biking, water exercise) rather than walking if neuropathy affects stability or use treadmill with safety handrails</li> <li>• Resistance training recommendations: <ul style="list-style-type: none"> <li>◦ Monitor discomfort in hands when using hand-held weights</li> <li>◦ Consider using dumbbells with soft/rubber coating, and/or wear padded gloves</li> <li>◦ Consider resistance machines over free weights (120)</li> </ul> </li> </ul>
Stem cell transplantation	<ul style="list-style-type: none"> <li>• Home-based exercise encouraged</li> <li>• A full recovery of the immune system recommended before return to gym facilities with the general public</li> <li>• Start with light intensity, short durations but high frequency and progress slowly (121)</li> <li>• Exercise volume (intensity and duration) should be adapted on a daily basis based on the individual's presentation</li> </ul>
Symptom clusters	<ul style="list-style-type: none"> <li>• Symptoms and side effects of cancer treatment rarely appear in isolation; rather, symptom clusters are the norm (i.e., fatigue, pain, sleep disturbance), especially during cancer treatment and in those with advanced disease (122).</li> <li>• Exercise professionals must be aware of this complexity and be prepared to refer clients/patients back to the medical team (i.e., rehabilitation or oncology physician, general practitioner, or nurse) for review and management of symptoms when safety concerns develop or when target symptom (e.g., fatigue) is not responding as expected.</li> </ul>
Sun safety	<ul style="list-style-type: none"> <li>• In addition to melanoma survivors (123), survivors of cancer at other primary sites may be at increased risk for secondary skin cancers (124)</li> <li>• Exercise professionals should recommend that cancer survivors engage in sun protective practices when exercising outdoors (125).</li> </ul>



insufficient evidence categories. In addition, the literature remains insufficient for further detailing prescriptions according to cancer type, timing of treatment, and/or types of treatment, whereas exercise prescriptions were rarely based on studies that directly compared varying FITT components, such as a head-to-head trial of low-intensity versus high-intensity training. Thus, as the evidence base continues to grow in other cancer sites and keeps pace with the evolution of cancer treatment, as well as trial designs broaden to include multiple treatment arms, the next generation of exercise prescriptions could have the specificity needed to move exercise oncology toward the same goal as precision oncology

where treatment is matched to the specific characteristics of a person's cancer.

Funding for the Roundtable was provided by: American College of Sports Medicine, American Cancer Society, American Academy of Physical Medicine and Rehabilitation, American Physical Therapy Association, Canadian Society for Exercise Physiology, Exercise and Sports Science Australia, German Union for Health Exercise and Exercise Therapy (DVGS), McMillan, Royal Dutch Society for Physical Therapy (KNGF), Sunflower Wellness.

No authors have conflicts of interest. The results of the present study do not constitute endorsement by ACSM.

Roundtable meeting attendees are listed in the following supplemental file (see Table, Meeting Attendees for the ACSM International Multidisciplinary Roundtable on Exercise and Cancer Prevention and Control, <http://links.lww.com/MSS/B789>).

## REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin*. 2019;69(1):7–34.
2. Bower JE. Management of cancer-related fatigue. *Clin Adv Hematol Oncol*. 2006;4(11):828–9.
3. Curigliano G, Cardinale D, Dent S, et al. Cardiotoxicity of anti-cancer treatments: epidemiology, detection, and management. *CA Cancer J Clin*. 2016;66(4):309–25.
4. Scott JM, Nilsen TS, Gupta D, Jones LW. Exercise therapy and cardiovascular toxicity in cancer. *Circulation*. 2018;137(11):1176–91.
5. Miller KD, Siegel RL, Lin CC, et al. Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin*. 2016;66(4):271–89.
6. Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc*. 2010;42(7):1409–26.
7. Department of Human and Health Services. Physical Activity Guidelines for Americans. Department of Health and Human Services 2008.
8. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology - Survivorship. 2018;Version 2.2018.
9. Cornie P, Atkinson M, Bucci L, et al. Clinical Oncology Society of Australia position statement on exercise in cancer care. *Med J Aust*. 2018;209(4):184–7.
10. Segal R, Zwaal C, Green E, Tomasone JR, Loblaw A, Petrella T. Exercise for people with cancer: a clinical practice guideline. *Curr Oncol*. 2017;24(1):40–6.
11. Rock CL, Doyle C, Demark-Wahnefried W, et al. Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin*. 2012;62:242–74.
12. Weggemans RM, Backx FJG, Borghouts L, et al. The 2017 Dutch Physical Activity Guidelines. *Int J Behav Nutr Phys Act*. 2018;15(1):58.
13. Mishra SI, Scherer RW, Geigle PM, et al. Exercise interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev*. 2012;(8):CD007566.
14. Mishra SI, Scherer RW, Snyder C, Geigle PM, Berlanstein DR, Topaloglu O. Exercise interventions on health-related quality of life for people with cancer during active treatment. *Cochrane Database Syst Rev*. 2012;8:CD008465.
15. Persoon S, Kersten MJ, van der Weiden K, et al. Effects of exercise in patients treated with stem cell transplantation for a hematologic malignancy: a systematic review and meta-analysis. *Cancer Treat Rev*. 2013;39(6):682–90.
16. Zhou Y, Zhu J, Gu Z, Yin X. Efficacy of exercise interventions in patients with acute leukemia: a meta-analysis. *PLoS One*. 2016; 11(7):e0159966.
17. Lahart IM, Metsios GS, Nevill AM, Carmichael AR. Physical activity for women with breast cancer after adjuvant therapy. *Cochrane Database Syst Rev*. 2018;1:CD011292.
18. Brown JC, Huedo-Medina TB, Pescatello LS, et al. The efficacy of exercise in reducing depressive symptoms among cancer survivors: a meta-analysis. *PLoS One*. 2012;7(1):e30955.
19. Craft LL, Vaniterson EH, Helenowski IB, Rademaker AW, Courneya KS. Exercise effects on depressive symptoms in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2012;21(1):3–19.
20. Puetz TW, Herring MP. Differential effects of exercise on cancer-related fatigue during and following treatment: a meta-analysis. *Am J Prev Med*. 2012;43(2):e1–24.
21. Tomlinson D, Diorio C, Beyene J, Sung L. Effect of exercise on cancer-related fatigue: a meta-analysis. *Am J Phys Med Rehabil*. 2014;93(8):675–86.
22. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Effects of supervised multimodal exercise interventions on cancer-related fatigue: systematic review and meta-analysis of randomized controlled trials. *Biomed Res Int*. 2015;2015:328636.
23. van Vulpen JK, Peeters PH, Velthuis MJ, van der Wall E, May AM. Effects of physical exercise during adjuvant breast cancer treatment on physical and psychosocial dimensions of cancer-related fatigue: a meta-analysis. *Maturitas*. 2016;85:104–11.
24. Brown JC, Huedo-Medina TB, Pescatello LS, Pescatello SM, Ferrer RA, Johnson BT. Efficacy of exercise interventions in modulating cancer-related fatigue among adult cancer survivors: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2011;20(1):123–33.
25. Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev*. 2012;11: CD006145.
26. Juvet LK, Thune I, Elvsaa IKO, et al. The effect of exercise on fatigue and physical functioning in breast cancer patients during and after treatment and at 6 months follow-up: a meta-analysis. *Breast*. 2017;33:166–77.
27. Keogh JW, MacLeod RD. Body composition, physical fitness, functional performance, quality of life, and fatigue benefits of exercise for prostate cancer patients: a systematic review. *J Pain Symptom Manage*. 2012;43(1):96–110.
28. Mustian KM, Alfano CM, Heckler C, et al. Comparison of pharmaceutical, psychological, and exercise treatments for cancer-related fatigue: a meta-analysis. *JAMA Oncol*. 2017;3(7):961–8.
29. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Supervised exercise reduces cancer-related fatigue: a systematic review. *J Physiother*. 2015;61(1):3–9.
30. McMillan EM, Newhouse IJ. Exercise is an effective treatment modality for reducing cancer-related fatigue and improving physical capacity in cancer patients and survivors: a meta-analysis. *Appl Physiol Nutr Metab*. 2011;36(6):892–903.
31. Buffart LM, Kalter J, Sweegers MG, et al. Effects and moderators of exercise on quality of life and physical function in patients with cancer: an individual patient data meta-analysis of 34 RCTs. *Cancer Treat Rev*. 2017;52:91–104.
32. Sweegers MG, Altenburg TM, Chinapaw MJ, et al. Which exercise prescriptions improve quality of life and physical function in patients with cancer during and following treatment? A systematic

- review and meta-analysis of randomised controlled trials. *Br J Sports Med*. 2018;52(8):505–13.
33. Erickson VS, Pearson ML, Ganz PA, Adams J, Kahn KL. Arm edema in breast cancer patients. *J Natl Cancer Inst*. 2001;93(2):96–111.
  34. Petrek JA, Pressman PI, Smith RA. Lymphedema: current issues in research and management. *CA Cancer J Clin*. 2000;50(5):292–307; quiz 308–11.
  35. Singh B, Disipio T, Peake J, Hayes SC. Systematic review and meta-analysis of the effects of exercise for those with cancer-related lymphedema. *Arch Phys Med Rehabil*. 2016;97(2):302–15.e13.
  36. Keilani M, Hasenoehrl T, Neubauer M, Crevenna R. Resistance exercise and secondary lymphedema in breast cancer survivors—a systematic review. *Support Care Cancer*. 2016;24(4):1907–16.
  37. Nelson NL. Breast cancer-related lymphedema and resistance exercise: a systematic review. *J Strength Cond Res*. 2016;30(9):2656–65.
  38. Ahmed RL, Thomas W, Yee D, Schmitz KH. Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol*. 2006;24(18):2765–72.
  39. Coumeya KS, Segal RJ, Mackey JR, et al. Effects of aerobic and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *J Clin Oncol*. 2007;25(28):4396–404.
  40. Schmitz KH, Ahmed RL, Troxel A, et al. Weight lifting in women with breast-cancer-related lymphedema. *N Engl J Med*. 2009;361(7):664–73.
  41. Schmitz KH, Ahmed RL, Troxel AB, et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *JAMA*. 2010;304(24):2699–705.
  42. Kilbreath SL, Refshauge KM, Beith JM, et al. Upper limb progressive resistance training and stretching exercises following surgery for early breast cancer: a randomized controlled trial. *Breast Cancer Res Treat*. 2012;133(2):667–76.
  43. Katz E, Dugan NL, Cohn JC, Chu C, Smith RG, Schmitz KH. Weight lifting in patients with lower-extremity lymphedema secondary to cancer: a pilot and feasibility study. *Arch Phys Med Rehabil*. 2010;91(7):1070–6.
  44. Swartz MC, Lewis ZH, Lyons EJ, et al. Effect of home- and community-based physical activity interventions on physical function among cancer survivors: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2017;98(8):1652–65.
  45. Dalla Via J, Daly RM, Fraser SF. The effect of exercise on bone mineral density in adult cancer survivors: a systematic review and meta-analysis. *Osteoporos Int*. 2018;29(2):287–303.
  46. Fornusek CP, Kilbreath SL. Exercise for improving bone health in women treated for stages I-III breast cancer: a systematic review and meta-analyses. *J Cancer Surviv*. 2017;11(5):525–41.
  47. Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR, American College of Sports Medicine. American College of Sports Medicine position stand: physical activity and bone health. *Med Sci Sports Exerc*. 2004;36(11):1985–96.
  48. Watson SL, Weeks BK, Weis LJ, Harding AT, Horan SA, Beck BR. High-intensity exercise did not cause vertebral fractures and improves thoracic kyphosis in postmenopausal women with low to very low bone mass: the LIFTMOR trial. *Osteoporos Int*. 2019;30(5):957–64.
  49. Chiu HY, Huang HC, Chen PY, Hou WH, Tsai PS. Walking improves sleep in individuals with cancer: a meta-analysis of randomized, controlled trials. *Oncol Nurs Forum*. 2015;42(2):E54–62.
  50. Mercier J, Savard J, Bernard P. Exercise interventions to improve sleep in cancer patients: a systematic review and meta-analysis. *Sleep Med Rev*. 2017;36:43–56.
  51. Rogers LQ, Coumeya KS, Oster RA, et al. Physical activity and sleep quality in breast cancer survivors: a randomized trial. *Med Sci Sports Exerc*. 2017;49(10):2009–15.
  52. Chen HM, Tsai CM, Wu YC, Lin KC, Lin CC. Effect of walking on circadian rhythms and sleep quality of patients with lung cancer: a randomised controlled trial. *Br J Cancer*. 2016;115(11):1304–12.
  53. Roveda E, Vitale JA, Bruno E, et al. Protective effect of aerobic physical activity on sleep behavior in breast cancer survivors. *Integr Cancer Ther*. 2017;16(1):21–31.
  54. Physical Activity Guidelines Advisory Committee. 2018 Physical Activity Guidelines Advisory Committee Scientific Report. In: *Department of Health and Human Services*. Washington, DC; 2018.
  55. Chen JJ, Wu PT, Middlekauff HR, Nguyen KL. Aerobic exercise in anthracycline-induced cardiotoxicity: a systematic review of current evidence and future directions. *Am J Physiol Heart Circ Physiol*. 2017;312(2):H213–22.
  56. Beaudry RI, Liang Y, Boyton ST, et al. Meta-analysis of exercise training on vascular endothelial function in cancer survivors. *Integr Cancer Ther*. 2018;17(2):192–9.
  57. Duregon F, Vendramin B, Bullo V, et al. Effects of exercise on cancer patients suffering chemotherapy-induced peripheral neuropathy undergoing treatment: a systematic review. *Crit Rev Oncol Hematol*. 2018;121:90–100.
  58. Campbell KL, Zdravec K, Bland KA, Chesley E, Wolf F, Janelsins M. The effect of exercise on cancer-related cognitive impairment and applications for physical therapy: a systematic review of randomized clinical trials. *Phys Ther*. doi.org/10.1093/ptj/pzz090.
  59. Northey JM, Cherbuin N, Pumpa KL, Smeed DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med*. 2018;52(3):154–60.
  60. Centres for Disease Control. *CDC Compendium of Effective Fall Interventions: What Works for Community-Dwelling Older Adults*. 3rd ed. 2015.
  61. Guirguis-Blake JM, Michael YL, Perdue LA, Coppola EL, Beil TL, Thompson JH. *Interventions to Prevent Falls in Community-Dwelling Older Adults: A Systematic Review for the US Preventive Services Task Force*. U.S. Preventive Services Task Force Evidence Syntheses, formerly Systematic Evidence Reviews. Rockville, MD; 2018.
  62. Winningham ML, MacVicar MG. The effect of aerobic exercise on patient reports of nausea. *Oncol Nurs Forum*. 1988;15(4):447–50.
  63. Irwin ML, Cartmel B, Gross CP, et al. Randomized exercise trial of aromatase inhibitor-induced arthralgia in breast cancer survivors. *J Clin Oncol*. 2015;33(10):1104–11.
  64. McNeely ML, Parliament MB, Seikaly H, et al. Effect of exercise on upper extremity pain and dysfunction in head and neck cancer survivors: a randomized controlled trial. *Cancer*. 2008;113(1):214–22.
  65. Yunfeng G, Weiyang H, Xueyang H, Yilong H, Xin G. Exercise overcome adverse effects among prostate cancer patients receiving androgen deprivation therapy: an update meta-analysis. *Medicine (Baltimore)*. 2017;96(27):e7368.
  66. Bourke L, Smith D, Steed L, et al. Exercise for men with prostate cancer: a systematic review and meta-analysis. *Eur Urol*. 2016;69(4):693–703.
  67. Cormie P, Chambers SK, Newton RU, et al. Improving sexual health in men with prostate cancer: randomised controlled trial of exercise and psychosexual therapies. *BMC Cancer*. 2014;14:199.
  68. Cormie P, Newton RU, Taaffe DR, et al. Exercise maintains sexual activity in men undergoing androgen suppression for prostate cancer: a randomized controlled trial. *Prostate Cancer Prostatic Dis*. 2013;16(2):170–5.
  69. Hojan K, Kwiatkowska-Borowczyk E, Leporowska E, et al. Physical exercise for functional capacity, blood immune function, fatigue, and quality of life in high-risk prostate cancer patients during radiotherapy: a prospective, randomized clinical study. *Eur J Phys Rehabil Med*. 2016;52(4):489–501.
  70. Stanton AM, Handy AB, Meston CM. The effects of exercise on sexual function in women. *Sex Med Rev*. 2018;6(4):548–57.

71. Gerbild H, Larsen CM, Graugaard C, Areskoung Josefsson K. Physical activity to improve erectile function: a systematic review of intervention studies. *Sex Med*. 2018;6(2):75–89.
72. Bland KA, Zdravec K, Landry T, Weller S, Meyers L, Campbell KL. Impact of exercise on chemotherapy completion rate: a systematic review of the evidence and recommendations for future exercise oncology research. *Crit Rev Oncol Hematol*. 2019;136:79–85.
73. Courneya KS. Physical activity and cancer survivorship: a simple framework for a complex field. *Exerc Sport Sci Rev*. 2014;42(3):102–9.
74. Winters-Stone KM, Neil SE, Campbell KL. Attention to principles of exercise training: a review of exercise studies for survivors of cancers other than breast. *Br J Sports Med*. 2014;48(12):987–95.
75. Neil-Sztramko SE, Medysky ME, Campbell KL, Bland KA, Winters-Stone KM. Attention to the principles of exercise training in exercise studies on prostate cancer survivors: a systematic review. *BMC Cancer*. 2019;19(1):321.
76. Neil-Sztramko SE, Winters-Stone KM, Bland KA, Campbell KL. Updated systematic review of exercise studies in breast cancer survivors: attention to the principles of exercise training. *Br J Sports Med*. 2019;53(8):504–12.
77. Courneya KS, McKenzie DC, Mackey JR, et al. Effects of exercise dose and type during breast cancer chemotherapy: multicenter randomized trial. *J Natl Cancer Inst*. 2013;105(23):1821–32.
78. van Waart H, Stuiver MM, van Harten WH, Geleijn E, Kieffers JM, Buffart LM, et al. Effect of low-intensity physical activity and Moderate- to high-intensity physical exercise during adjuvant chemotherapy on physical fitness, fatigue, and chemotherapy completion rates: Results of the PACES randomized clinical trial. *J Clin Oncol*. 2015;33(17):1918–27.
79. Danhauer SC, Addington EL, Cohen L, et al. Yoga for symptom management in oncology: a review of the evidence base and future directions for research. *Cancer*. 2019;125(12):1979–89.
80. McKenzie DC. Abreast in a boat—a race against breast cancer. *CMAJ*. 1998;159(4):376–8.
81. McDonough MH, Patterson MC, Weisenbach BB, Ullrich-French S, Sabiston CM. The difference is more than floating: factors affecting breast cancer survivors' decisions to join and maintain participation in dragon boat teams and support groups. *Disabil Rehabil*. 2019;41:1788–96.
82. Uth J, Hornstrup T, Christensen JF, et al. Efficacy of recreational football on bone health, body composition, and physical functioning in men with prostate cancer undergoing androgen deprivation therapy: 32-week follow-up of the FC prostate randomised controlled trial. *Osteoporos Int*. 2016;27(4):1507–18.
83. Crawford JJ, Vallance JK, Holt NL, Bell GJ, Steed H, Courneya KS. A pilot randomized, controlled trial of a Wall climbing intervention for gynecologic cancer survivors. *Oncol Nurs Forum*. 2017;44(1):77–86.
84. Ng AV, Cybulski AN, Engel AA, et al. Triathlon training for women breast cancer survivors: feasibility and initial efficacy. *Support Care Cancer*. 2017;25(5):1465–73.
85. Dolan LB, Campbell K, Gelmon K, Neil-Sztramko S, Holmes D, McKenzie DC. Interval versus continuous aerobic exercise training in breast cancer survivors—a pilot RCT. *Support Care Cancer*. 2016;24(1):119–27.
86. Riebe D, Ehrman JK, Liguori G, Magal M, editors. *ACSM's Guidelines for Exercise Testing and Prescription 10th ed*. Philadelphia, PA: Wolters Kluwer; 2018.
87. McNeely ML, Courneya KS. Exercise programs for cancer-related fatigue: evidence and clinical guidelines. *J Natl Compr Canc Netw*. 2010;8(8):945–53.
88. Galvao DA, Taaffe DR, Spry N, et al. Exercise preserves physical function in prostate cancer patients with bone metastases. *Med Sci Sports Exerc*. 2018;50(3):393–9.
89. Panel on Prevention of Falls in Older Persons, American Geriatrics Society and British Geriatrics Society. Summary of the updated American Geriatrics Society/British Geriatrics Society Clinical Practice Guideline for Prevention of Falls in Older Persons. *J Am Geriatr Soc*. 2011;59(1):148–57.
90. Patnaik JL, Byers T, DiGiuseppi C, Dabelea D, Denberg TD. Cardiovascular disease competes with breast cancer as the leading cause of death for older females diagnosed with breast cancer: a retrospective cohort study. *Breast Cancer Res*. 2011;13(3):R64.
91. Riebe D, Franklin BA, Thompson PD, et al. Updating ACSM's recommendations for exercise preparticipation health screening. *Med Sci Sports Exerc*. 2015;47(11):2473–9.
92. Scott JM, Zabor EC, Schwitzer E, et al. Efficacy of exercise therapy on cardiorespiratory fitness in patients with cancer: a systematic review and meta-analysis. *J Clin Oncol*. 2018;36(22):2297–305.
93. Fuller JT, Hartland MC, Maloney LT, Davison K. Therapeutic effects of aerobic and resistance exercises for cancer survivors: a systematic review of meta-analyses of clinical trials. *Br J Sports Med*. 2018;52(20):1311.
94. Cheema BS, Kilbreath SL, Fahey PP, Delaney GP, Atlantis E. Safety and efficacy of progressive resistance training in breast cancer: a systematic review and meta-analysis. *Breast Cancer Res Treat*. 2014;148(2):249–68.
95. Padilha CS, Marinello PC, Galvao DA, et al. Evaluation of resistance training to improve muscular strength and body composition in cancer patients undergoing neoadjuvant and adjuvant therapy: a meta-analysis. *J Cancer Surviv*. 2017;11(3):339–49.
96. Sweegers MG, Altenburg TM, Brug J, et al. Effects and moderators of exercise on muscle strength, muscle function and aerobic fitness in patients with cancer: a meta-analysis of individual patient data. *Br J Sports Med*. 2019;53(13):812.
97. Buffart LM, Sweegers MG, May AM, et al. Targeting exercise interventions to patients with cancer in need: an individual patient data meta-analysis. *J Natl Cancer Inst*. 2018;110(11):1190–200.
98. Dolan LB, Gelmon K, Courneya KS, et al. Hemoglobin and aerobic fitness changes with supervised exercise training in breast cancer patients receiving chemotherapy. *Cancer Epidemiol Biomarkers Prev*. 2010;19(11):2826–32.
99. Lakoski SG, Eves ND, Douglas PS, Jones LW. Exercise rehabilitation in patients with cancer. *Nat Rev Clin Oncol*. 2012;9(5):288–96.
100. Galvao DA, Newton RU, Taaffe DR, Spry N. Can exercise ameliorate the increased risk of cardiovascular disease and diabetes associated with ADT? *Nat Clin Pract Urol*. 2008;5(6):306–7.
101. Galvao DA, Taaffe DR, Spry N, Joseph D, Turner D, Newton RU. Reduced muscle strength and functional performance in men with prostate cancer undergoing androgen suppression: a comprehensive cross-sectional investigation. *Prostate Cancer Prostatic Dis*. 2009;12(2):198–203.
102. Gardner JR, Livingston PM, Fraser SF. Effects of exercise on treatment-related adverse effects for patients with prostate cancer receiving androgen-deprivation therapy: a systematic review. *J Clin Oncol*. 2014;32(4):335–46.
103. Winters-Stone KM, Dobek JC, Bennett JA, et al. Resistance training reduces disability in prostate cancer survivors on androgen deprivation therapy: evidence from a randomized controlled trial. *Arch Phys Med Rehabil*. 2015;96(1):7–14.
104. Baracos VE, Martin L, Korc M, Guttridge DC, Fearon KCH. Cancer-associated cachexia. *Nat Rev Dis Primers*. 2018;4:17105.
105. Demark-Wahnefried W, Schmitz KH, Alfano CM, et al. Weight management and physical activity throughout the cancer care continuum. *CA Cancer J Clin*. 2018;68(1):64–89.
106. World Cancer Research Fund/American Institute for Cancer Research. Diet, physical activity and cancer: a global perspective. 2018. <https://www.wcrf.org/dietandcancer>. Access May 1, 2019.
107. Arends J, Baracos V, Bertz H, et al. ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr*. 2017;36(5):1187–96.
108. Arends J, Bachmann P, Baracos V, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr*. 2017;36(1):11–48.



109. American College of Sports Medicine, Armstrong LE, Casa DJ, Millard-Stafford M, et al. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc.* 2007;39(3):556–72.
110. Patel AV, Friedenreich CM, Moore SC, et al. American College of Sports Medicine Roundtable report on physical activity, sedentary behavior, and cancer prevention and control. *Med Sci Sports Exerc.* 2019;51(11):2391–402.
111. Rief H, Petersen LC, Omlor G, et al. The effect of resistance training during radiotherapy on spinal bone metastases in cancer patients—a randomized trial. *Radiother Oncol.* 2014;112(1):133–9.
112. Frost HM. Should future risk-of-fracture analyses include another major risk factor? The case for falls. *J Clin Densitom.* 2001;4(4):381–3.
113. Paskett ED, Dean JA, Oliveri JM, Harrop JP. Cancer-related lymphedema risk factors, diagnosis, treatment, and impact: a review. *J Clin Oncol.* 2012;30(30):3726–33.
114. Clough-Gorr KM, Stuck AE, Thwin SS, Silliman RA. Older breast cancer survivors: geriatric assessment domains are associated with poor tolerance of treatment adverse effects and predict mortality over 7 years of follow-up. *J Clin Oncol.* 2010;28(3):380–6.
115. Klepin HD, Geiger AM, Tooze JA, et al. Physical performance and subsequent disability and survival in older adults with malignancy: results from the health, aging and body composition study. *J Am Geriatr Soc.* 2010;58(1):76–82.
116. Maccormick RE. Possible acceleration of aging by adjuvant chemotherapy: a cause of early onset frailty? *Med Hypotheses.* 2006;67(2):212–5.
117. American College of Sports Medicine, Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, et al. American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc.* 2009;41(7):1510–30.
118. Association of Stoma Care Nurses UK. ACSN Stoma Care National Clinical Guidelines. 2016. <http://ascnuk.com/wp-content/uploads/2016/03/ASCN-Clinical-Guidelines-Final-25-April-compressed-11-10-38.pdf> (Access Date May 1, 2019).
119. Esper P. Symptom clusters in individuals living with advanced cancer. *Semin Oncol Nurs.* 2010;26(3):168–74.
120. Streckmann F, Zopf EM, Lehmann HC, et al. Exercise intervention studies in patients with peripheral neuropathy: a systematic review. *Sports Med.* 2014;44(9):1289–304.
121. Wiskemann J, Huber G. Physical exercise as adjuvant therapy for patients undergoing hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2008;41(4):321–9.
122. Ward Sullivan C, Leutwyler H, Dunn LB, Miaskowski C. A review of the literature on symptom clusters in studies that included oncology patients receiving primary or adjuvant chemotherapy. *J Clin Nurs.* 2018;27(3-4):516–45.
123. Tucker MA, Elder DE, Curry M, et al. Risks of melanoma and other cancers in melanoma-prone families over 4 decades. *J Invest Dermatol.* 2018;138(7):1620–6.
124. Donin N, Filson C, Drakaki A, et al. Risk of second primary malignancies among cancer survivors in the United States, 1992 through 2008. *Cancer.* 2016;122(19):3075–86.
125. Lau SC, Chen L, Cheung WY. Protective skin care behaviors in cancer survivors. *Curr Oncol.* 2014;21(4):e531–40.
126. Schmitz KH, al. E. Exercise Is Medicine in Oncology: Moving Through Cancer. *CA Cancer J Clin.* In press.